



TEXTURED

SHOW FEEDS

0776

PRODUCT DATA

DIG. ENERGY	1465.48	Kcal/lb
ME	1.38	Mcal/lb
NE MAINTENANCE ..	0.79	Mcal/lb
NE GAIN	0.50	Mcal/lb
TDN	73.16	%
PROTEIN	12.05	%
DIG. PROTEIN	10.39	%
NPN	1.72	%
LYSINE	0.44	%
FAT	2.12	%
FIBER	13.76	%
ASH	3.85	%
CELL WALLS	30.17	%
ADF	20.66	%
CELLULOSE	12.67	%
LIGNIN	5.74	%
CALCIUM	0.61	%
CHLORINE	1.04	%
COBALT	0.32	mg/lb
COPPER	8.72	mg/lb
IRON	61.64	mg/lb
IODINE	0.27	mg/lb
MAGNESIUM	0.23	%
MANGANESE	32.28	mg/lb
PHOSPHORUS	0.31	%
POTASSIUM	1.14	%
SODIUM	0.44	%
SELENIUM	0.04	mg/lb
SULFUR	0.22	%
ZINC	34.45	mg/lb
RIBOFLAVIN	0.76	mg/lb
THIAMINE	0.92	mg/lb
NIACIN	8.89	mg/lb
PAN ACID	4.31	mg/lb
CHOLINE	333.35	mg/lb
PYRIDOXINE	3.53	mg/lb
FOLIC ACID	0.16	mg/lb
VITAMIN B12	0.00	mcg/lb
VITAMIN E	6.97	IU/lb
VITAMIN A	5.50	KIU/lb
VITAMIN D	1.23	KIU/lb

TFP NUTRITION
NACOGDOCHES, TEXAS

BULL & HEIFER DEVELOPER


A medicated, cottonseed hull and corn based ration used by many cattlemen raising purebred bulls and heifers for replacement or sale.

FEATURES

- * 1465 kcal/lb
- * 12% protein
- * Medicated with Bovatec®
- * 21% cottonseed hulls
- * 45% corn
- * Contains soybean hulls

BENEFITS

- * Corn provides optimum finish on animals.
- * Medication improves rate of weight gain.
- * Excellent for fitting cattle for show.
- * Cottonseed hulls and soybean hulls provide high quality roughage.



FOR BEEF CATTLE ON PASTURE

For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Lasalocid 27 grams/ton

GUARANTEED ANALYSIS

Crude Protein, minimum	12.00%
(This includes not more than 1.72% equivalent crude protein from non-protein nitrogen.)	
Crude Fat, minimum	2.00%
Crude Fiber, maximum	18.00%
Calcium (Ca), minimum	0.45%
Calcium (Ca), maximum	0.75%
Phosphorus (P), minimum	0.30%
Salt (NaCl), minimum	0.75%
Salt (NaCl), maximum	1.25%
Potassium (K), minimum	0.50%
Vitamin A, minimum	5,000 IU/Lb
Vitamin D3, minimum	1,230 IU/Lb

FEEDING DIRECTIONS

Feed continuously to beef cattle on pasture at the rate of 4.5 to 15 pounds per head per day. This will provide a minimum of 60 mg and a maximum of 200 mg of lasalocid per head per day. Provide a source of fresh, clean water at all times.

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to processed for veal.

CAUTION: The safety of Lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.