

TEXTURED

Medicated

CATTLE FEEDS

0795

PRODUCT DATA

| DIG. ENERGY | 1462.20 | Kcal/lb |
|----------------|---------|---------|
| ME | 1.37 | Mcal/lb |
| NE MAINTENANCE | 0.78 | Mcal/lb |
| NE GAIN | 0.50 | Mcal/lb |
| TDN | 72.94 | % |
| PROTEIN | 12.12 | % |
| DIG. PROTEIN | 10.19 | % |
| NPN | 0.00 | % |
| LYSINE | 0.52 | % |
| FAT | 5.00 | % |
| FIBER | 12.81 | % |
| ASH | 4.04 | % |
| CELL WALLS | 29.55 | % |
| ADF | 19.35 | % |
| CELLULOSE | 11.28 | % |
| LIGNIN | 6.21 | % |
| CALCIUM | 0.71 | % |
| CHLORINE | 91.88 | % |
| COBALT | 0.33 | mg/lb |
| COPPER | 10.09 | mg/lb |
| IRON | 87.95 | mg/lb |
| IODINE | 0.27 | mg/lb |
| MAGNESIUM | 0.26 | % |
| MANGANESE | 43.98 | mg/lb |
| PHOSPHORUS | 0.46 | % |
| POTASSIUM | 1.15 | % |
| SODIUM | 0.26 | % |
| SELENIUM | 0.05 | mg/lb |
| SULFUR | 0.26 | % |
| ZINC | 37.12 | mg/lb |
| RIBOFLAVIN | 0.89 | mg/lb |
| THIAMINE | 1.26 | mg/lb |
| NIACIN | 9.70 | mg/lb |
| PAN ACID | 5.09 | mg/lb |
| CHOLINE | | mg/lb |
| PYRIDOXINE | 3.16 | mg/lb |
| FOLIC ACID | 0.22 | mg/lb |
| VITAMIN B12 | 0.01 | mcg/lb |
| VITAMIN E | 6.86 | IU/lb |
| VITAMIN A | 5.43 | KIU/lb |
| MITALAINED | 4.00 | 120110 |

TEXAS FARM PRODUCTS COMPANY NACOGDOCHES, TEXAS

FITTER'S CHOICE CALF FINISHER


A high energy, corn, cottonseed hull, and oat based sweet feed formulated to provide an optimum finish to show calves.

FEATURES

- 1462 Kcal/lb
- 12% Protein
- Flaked Corn
- Crimped Oats
- Cottonseed Hulls
- Roasted Soybeans
- Rumen Bypass Fat
- Contains Boyatec®
- Fortified with **Probiotics**

BENEFITS

- * High corn content and roasted soybeans provides optimum finish on animals.
- Cottonseed hulls provide high quality roughage.
- Molasses improves palatability to ensure consistent feed consumption.
- * Bovatec® improve feed efficiency.
- * Beneficial microbes, yeast culture, and enzymes contained in the probiotics improves digestion, performance, and overall health.



For improved feed efficiency in cattle fed in confinement for slaughter.

ACTIVE DRUG INGREDIENT

| Lasalocid | | 27 | grams/ton |
|-----------|--|----|-----------|
|-----------|--|----|-----------|

| GUARANTEED ANALYSIS | |
|-------------------------|-------------|
| Crude Protein, minimum | 12.00% |
| Crude Fat, minimum | 5.00% |
| Crude Fiber, maximum | 13.00% |
| Calcium (Ca), minimum | 0.60% |
| Calcium (Ca), maximum | 1.10% |
| Phosphorus (P), minimum | 0.45% |
| Salt (NaCl), minimum | 0.25% |
| Salt (NaCl), maximum | 0.75% |
| Potassium (K), minimum | 0.80% |
| Vitamin A, minimum | 5,000 IU/LB |
| | |

INGREDIENTS

Flaked yellow corn, cottonseed hulls (21%), cane molasses, crimped oats, roasted soybeans, cottonseed meal, rice bran, soybean meal, pearled barley, calcium carbonate, calcium salts of long-chain fatty acids, dicalcium phosphate, salt, yeast culture, dried Lactobacillus acidophilus fermentation product. dried Enterococcus faecium fermentation product, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, ethylenediamine dihydroiodide, cobalt carbonate,

FEEDING DIRECTIONS

Feed continuously at the rate of 7.4 lb/head/day to 26.7 lb/head/day to show calves beginning approximately 90 to 140 days before show date. This will provide a minimum of 100 mg/ head/day of Lasalocid and a maximum of 360 mg/head/day. The length of finishing period depends on the the breed of calf, frame size, and degree of finish desired. As a general rule, begin feeding Fitter's Choice Calf Finisher to large frame calves at approximately 700 pound of body weight. Early maturing, small framed calves should be fed this ration beginning at approximately 800 pounds of body weight. Provide a source of fresh, clean water at all times throughout the feeding program.

WARNING: A withdrawal period has not been established for this product in preruminating cattle. Do not use in calves to be processed for veal.

CAUTION: The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. Changes in the feeding rate should be made gradually. Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.