

For Improved Digestion. Health, Appearance, and Overall Performance

HORSE FEEDS

1/8" PELLETS

1098

PRODUCT DATA DIG. ENERGY 1401.56 Kcal/lb

PROTEIN 16.10 %

LYSINE 0.84 %

FAT 3.40 %

FIBER 9.96 %

NDF 19.48 %

ADF 15.76 %

ASH 7.03 %

PHOSPHORUS 0.53 %

%

%

%

%

CALCIUM 0.84

MAGNESIUM 0.40

POTASSIUM 1.49

SODIUM 0.29

SULFUR 0.23 %

COPPER 10.97 mg/lb

IODINE 0.43 mg/lb

IRON 76.62 mg/lb

MANGANESE 52.60 mg/lb

SELENIUM 0.23 mg/lb

ZINC 33.58 mg/lb

COBALT 0.18 mg/lb

CAROTENE 6.11 mg/lb VITAMIN A 5.45 KIU/lb

A pelletized, vitamin fortified ration formulated especially for brood mares and foals.

BENEFITS

* Extra nutrition for foals.

and nutrient utilization.

feed conversion.

* Excellent for breeding stallions.

FEATURES

- 16% protein
- * High energy-1401 Kcal/lb * 10% Alfalfa
- With vitamins A. D. & E

SUPER GLO® 4

- With a Probiotic that contains:
- 1. Encapsulated microbes
- Streptococcus faecium &

GUARANTEED ANALYSIS

Crude Protein, minimum	16.00%
Crude Fat, minimum	. 3.00%
Crude Fiber, maximum	10.00%
Calcium (Ca), minimum	
Calcium (Ca), maximum	
Phosphorus (P), minimum	
Copper (Cu), minimum	
Selenium (Se), minimum 0	
Zinc (Zn), minimum	
Vitamin A, minimum 3,00	
Vitamin D3, minimum 3	
Vitamin E, minimum	36 IU/LB

* Added microbes colonize intestine with

* Mares remain in top condition when nursing.

* Yeast culture improves palatability, digestion,

protective microflora resulting in better health.

Added enzymes improve digestion of proteins,

starches, and cellulose resulting in improved

INGREDIENTS Grain products, processed grain by-products, plant protein products, roughage products (6%), forage products, can molasses, calcium carbonate, salt, *Streptococcus faecium* fermentation product, dried *Lactobacillus acidophilus* fermentation product, dried Aspergillus oryzae fermentation extract, dried Aspergillus niger fermentation extract, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, vitamin B12 supplement, riboflavin supplement, niacin supplement, calcium pantothenate, choline chloride, folic acid, thiamine mononitrate, pyridoxine hydrochloride, biotin, sodium selenite, zinc sulfate, ferrous sulfate, manganese sulfate, copper sulfate, ethylenediamine dihydroiodide.

FEEDING DIRECTIONS

The recommended feeding rate for SUPER GLO® 4 is shown below. Do not feed free choice except as directed. Provide plenty of good quality hay or pasture along with a source of fresh, clean water. If you feed by dry measure, periodically weigh the feed to ensure that the proper amount is being fed. Observe the condition of your horse over a period of time. The recommended feeding rates may be increased or decreased by 10% to keep your horse in the desired condition. Do not exceed 10% above the recommended rate.

Changes in feed or feeding rate should be made gradually over several days. Changes in feeding rate should not exceed 1 pound per day. Feed at regular times at least twice daily. If horse is hot, excited, or ill, delay feeding. Do not allow horse to over-consume or eat rapidly.

An effective parasite control program is essential to your horse's health. Observe your horse's condition regularly. Consult a veterinarian if any problems arise.

NURSING FOALS

Feed SUPER GLO® 4 free choice as a creep feed. Provide access to guality hav or pasture. Do not allow mares to consume this feed free choice. Do not allow foals to consume more than 2 lbs. of this feed per 100 pounds of body weight per day.

WEANED FOALS WEIGHT OF FOAL Hay/Pasture (min. lb/day) SUPER GLO® 4 (min.lbs./day)	300 4 7	400 5 8	500 6 8 ¹ /2	600 7 8 ¹ /2	
BREEDING HORSES WEIGHT OF HORSE	600	800	1000	1200	1400
Hay/Pasture (min. lb/day)	7	8	10	12	14
SUPER GLO® 4 (min. lb/day)					
Mares/Gestation (last 90 days)	5	6	7	8	9
Lactating Mares (first 90 days)*	10	13	15	18	20
Breeding Stallions	6	7	8	9	10

*Gradually reduce feeding rate after 90 days to maintain desired condition.



407210

Lactobacillus acidophilus 2. Yeast culture 3. Enzymes - Protease, Amylase, & Cellulase

Crude Fat, minimum 3.00%
Crude Fiber, maximum 10.00%
Calcium (Ca), minimum 0.50%
Calcium (Ca), maximum 1.00%
Phosphorus (P), minimum 0.45%
Copper (Cu), minimum 20 ppm
Selenium (Se), minimum 0.20 ppm
Zinc (Zn), minimum 50 ppm
Vitamin Á, minimum 3,000 IÚ/LB
Vitamin D3, minimum 375 IU/LB
Vitamin E, minimum

VITAMIN D 0.37 KIU/lb VITAMIN E 45.26 IU/lb BIOTIN 0.34 mg/lb FOLATE 0.86 mg/lb NIACIN 42.34 mg/lb PANTOTHENIC ACID 10.84 mg/lb RIBOFLAVIN 3.52 mg/lb THIAMINE 4.95 mg/lb PYRIDOXINE 4.51 mg/lb VITAMIN B12 10.00 mcg/lb

TFP NUTRITION NACOGDOCHES, TEXAS

CAUTION: Feed is perishable. Feed should be stored in a well-ventilated, dry area that is protected from rodents and insects. Do not feed moldy or insect-infested feed as it may cause illness or death.