



POULTRY FEEDS

1250

PRODUCT DATA

ME	1485.10 Kcal/lb
PROTEIN	23.05%
ARGININE	1.47%
GLYCINE	1.02%
SERINE	1.24%
HISTIDINE	0.58%
ISOLEUCINE	1.10%
LEUCINE	2.43%
LYSINE	1.21%
METHIONINE	0.58%
CYSTINE	0.37%
PHENYLALANINE	1.14%
TYROSINE	1.02%
THREONINE	0.91%
TRYPTOPHAN	0.26%
VALINE	1.26%
FAT	6.04%
LINOLEIC ACID	1.48%
FIBER	2.68%
CALCIUM	1.08%
TOTAL PHOS.	0.72%
AVAILABLE PHOS.	0.48%
POTASSIUM	0.76%
CHLORINE	0.20%
COBALT	0.24mg/lb
IODINE	0.58mg/lb
IRON	174.48 mg/lb
MAGNESIUM	0.19%
MANGANESE	41.53mg/lb
SODIUM	0.13%
SULFUR	0.22%
COPPER	5.16mg/lb
SELENIUM	0.19mg/lb
ZINC	41.83mg/lb
BIOTIN	0.14mg/lb
CHOLINE	663.57 mg/lb
FOLIC ACID	0.87mg/lb
NIACIN	29.71mg/lb
PANTOTHENIC ACID	8.23mg/lb
PYRIDOXINE	4.32mg/lb
RIBOFLAVIN	3.03mg/lb
THIAMINE	2.65mg/lb
VITAMIN A	3.63KIU/lb
VITAMIN D3	1.25KIU/lb
VITAMIN B12	6.18mcg/lb
VITAMIN E	16.77IU/lb

TEXAS FARM PRODUCTS COMPANY
NACOGDOCHES, TX

20% BROILER GROW & SHOW

A high-tech, state-of-the-art ration designed to produce heavy weight broilers for competition or market.

FEATURES

- * 23% protein
- * High energy - 1458 Kcal/lb.
- * Added fat
- * Crumblized
- * Added methionine and lysine
- * 7.5% corn gluten meal
- * Contains Amprolium as a coccidiostat
- * Fortified with special vitamins and minerals
- * Fortified with Probiotics

BENEFITS

- * Produces golden pigmentation.
- * Supplies balanced quantities of protein, amino acids, vitamins, minerals and energy to promote rapid growth.
- * Aids in the prevention of coccidiosis.
- * Added microbes colonize intestine with protective microflora resulting in better health.
- * Yeast culture improves digestion and nutrient utilization.
- * Added enzymes improve digestion of proteins, starches, and cellulose resulting in improved feed conversion.

As an aid in the prevention of coccidiosis.

ACTIVE DRUG INGREDIENTS

Amprolium	0.0125%
-----------	---------

GUARANTEED ANALYSIS

Crude Protein, minimum	23.00%
Lysine, minimum	1.10%
Methionine, minimum	0.50%
Crude Fat, minimum	6.00%
Crude Fiber, maximum	3.50%
Calcium (Ca), minimum	0.95%
Calcium (Ca), maximum	1.20%
Phosphorus (P), minimum	0.60%
Salt (NaCl), minimum	0.20%
Salt (NaCl), maximum	0.40%

FEEDING DIRECTIONS

DIRECTIONS FOR USE

Feed as the sole ration to broiler chicks from day of hatch to time of show. Keep feeders clean and provide a source of fresh, clean water at all times. Follow a strict sanitation program.

WARNING: Use as a sole source of Amprolium.

CAUTION: Feed should be stored in a well-ventilated, dry area that is protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.

