



GAME BIRD FEEDS

1271

PRODUCT DATA

ME	890.93	Kcal/lb
PROTEIN	20.09	%
ARGININE	1.36	%
GLYCINE	1.13	%
SERINE	0.99	%
HISTIDINE	0.47	%
ISOLEUCINE	0.88	%
LEUCINE	1.53	%
LYSINE	1.06	%
METHIONINE	0.45	%
CYSTINE	0.28	%
PHENYLALANINE	0.80	%
TYROSINE	0.70	%
THREONINE	0.69	%
TRYPTOPHAN	0.23	%
VALINE	1.01	%
FAT	3.08	%
LINOLEIC ACID	0.92	%
FIBER	8.03	%
CALCIUM	3.26	%
TOTAL PHOS.	0.71	%
AVAILABLE PHOS.	0.40	%
POTASSIUM	1.05	%
CHLORINE	0.36	%
COBALT	0.23	mg/lb
IODINE	0.66	mg/lb
IRON	43.91	mg/lb
MAGNESIUM	0.39	%
MANGANESE	74.44	mg/lb
SODIUM	0.28	%
SULFUR	0.22	%
COPPER	4.67	mg/lb
SELENIUM	0.19	mg/lb
ZINC	49.97	mg/lb
BIOTIN	0.13	mg/lb
CHOLINE	698.63	mg/lb
FOLIC ACID	0.63	mg/lb
NIACIN	30.84	mg/lb
PAN. ACID	9.80	mg/lb
PYRIDOXINE	2.77	mg/lb
RIBOFLAVIN	2.95	mg/lb
THIAMINE	3.11	mg/lb
VITAMIN A	3.69	KIU/lb
VITAMIN D3	1.25	KIU/lb
VITAMIN B12	4.78	mcg/lb
VITAMIN E	9.38	IU/lb

TFP NUTRITION
NACOGDOCHES, TX

GAME BIRD BREEDER-LAYER

A complete, crumbled ration designed specifically to meet the extra nutritional requirements for laying pheasants and quail.

FEATURES

- * 890.93 kcal/lb
- * 20% protein
- * 4:1 calcium to phosphorus ratio
- * Fortified with vitamins and minerals

BENEFITS

- * Increased egg production
- * Harder egg shells
- * Supplies all the protein, vitamins, minerals, and energy needed by laying birds

GUARANTEED ANALYSIS

Crude Protein, minimum	20.00%
Lysine, minimum	1.00%
Methionine, minimum	0.45%
Crude Fat, minimum	3.00%
Crude Fiber, maximum	8.00%
Calcium (Ca), minimum	2.50%
Calcium (Ca), maximum	3.50%
Phosphorus (P), minimum	0.70%
Salt (NaCl), minimum	0.25%
Salt (NaCl), maximum	0.75%

INGREDIENTS

Grain products, plant protein products, processed grain by-products, forage products, roughage products (10%), calcium carbonate, cane molasses, DL-methionine, salt, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, vitamin B12 supplement, riboflavin supplement, niacin supplement, calcium pantothenate, choline chloride, menadione sodium bisulfite complex, folic acid, pyridoxine hydrochloride, thiamine mononitrate, sodium selenite, biotin, manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, calcium iodate, cobalt sulfate.

FEEDING DIRECTIONS

Begin feeding free choice to pheasants and quail approximately one month before time of first egg and continue throughout the time of egg production. Provide a source of fresh, clean water at all times.

CAUTION: Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to birds as it may cause illness or death.

