

GAME BIRD FEEDS

1272

PRODUCT DATA

PRODUCT DATA		
ME	861.55	Kcal/lb
PROTEIN	19.06	%
ARGININE	1.29	%
GLYCINE	0.98	%
SERINE	0.97	%
HISTIDINE	0.46	%
ISOLEUCINE		%
LEUCINE		%
LYSINE	1.02	%
METHIONINE		%
CYSTINE	0.27	%
PHENYLALANINE		%
TYROSINE	0.69	%
THREONINE	0.67	%
TRYPTOPHAN		%
VALINE	0.98	%
FAT	2.77	%
LINOLEIC ACID	0.88	%
FIBER		%
CALCIUM	0.96	%
TOTAL PHOS	0./4	%
AVAILABLE PHOS.		%
POTASSIUM		%
CHLORINE		%
COBALTIODINE	0.24	mg/lb
		mg/lb
IRON	/0.42	mg/lb %
MAGNESIUM MANGANESE	00.60	
	80.62	mg/lb %
SODIUMSULFUR		% %
COPPER		mg/lb
SELENIUM		mg/lb
ZINC		mg/lb
BIOTIN		mg/lb
CHOLINE		mg/lb
FOLIC ACID		mg/lb
NIACIN		mg/lb
PAN. ACID		mg/lb
PYRIDOXINE		mg/lb
RIBOFLAVIN	2 90	mg/lb
THIAMINE		mg/lb
VITAMIN A		KIU/lb
VITAMIN D3		KIU/lb
VITAMIN B12		mcg/lb
VITAMIN E		IU/lb
		TOTIO

TEXAS FARM PRODUCTS COMPANY NACOGDOCHES, TX

GAME BIRD FLIGHT CONDITIONER

A complete feed for pheasants and quail raised for breeding or release on game preserves.

FEATURES

- * 19% protein
- * 861 Kcal/lb
- Vitamin and mineral fortified
- * 1/8 inch diameter pellets

BENEFITS

- * Protein to energy ratio promotes lean birds suitable for release.
- * Prevents excessive weight gains in young birds to be kept for breeding.
- * Convenient -- feed as the sole ration.

POR GAME BIRD FLIGHT CONDITIONER FOR GAME BIRD FLIGHT CONDITIONER FOR GAME BIRDS A TURKEYS

GUARANTEED ANALYSIS

Crude Protein, minimum	19.00%
Crude Fat, minimum	2.00%
Crude Fiber, maximum	12.00%
Calcium (Ca), minimum	0.80%
Calcium (Ca), maximum	1.10%
Phosphorus (P), minimum	0.70%
Salt (NaCl), minimum	0.25%
Salt (NaCl), maximum	0.75%

INGREDIENTS

Grain products, plant protein products, processed grain by-products, roughage products (20%), forage products, cane molasses, dicalcium phosphate, calcium carbonate, salt, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, vitamin B12 supplement, riboflavin supplement, niacin supplement, calcium pantothenate, choline chloride, menadione sodium bisulfite complex, folic acid, pyridoxine hydrochloride, thiamine mononitrate, sodium selenite, biotin, manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, calcium iodate, cobalt sulfate.

FEEDING DIRECTIONS

Feed free choice from 8 to 18 weeks of age to pheasants and quail grown for breeding or release on the game preserve. Provide a source of fresh, clean water at all times.

CAUTION: Changes in the feeding rate should be made gradually. Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.