



For Improved Digestion,
Health, Appearance, and
Overall Performance

SHOW FEEDS

1673

PRODUCT DATA

DIG. ENERGY	1351.01	Kcal/lb
ME	1.06	Mcal/lb
NE MAINTENANCE ..	0.76	Mcal/lb
NE GAIN	0.51	Mcal/lb
TDN	68.33	%
PROTEIN	16.00	%
DIG. PROTEIN	14.12	%
NPN	1.00	%
CELLULOSE	6.41	%
FAT	4.59	%
FIBER	7.93	%
LIGNIN	2.31	%
CALCIUM	1.18	%
CHLORINE	0.52	%
COBALT	0.27	mg/lb
COPPER	4.68	mb/lb
IODINE	0.22	mg/lb
IRON	60.93	mg/lb
MAGNESIUM	0.20	%
MANGANESE	39.51	mg/lb
PHOSPHORUS	0.48	%
POTASSIUM	0.99	%
SELENIUM	0.10	mg/lb
SODIUM	0.33	%
SULFUR	0.20	%
ZINC	35.22	mg/lb
CAROTENE	0.33	mg/lb
VITAMIN A	5.08	KIU/lb
VITAMIN D	1.23	KIU/lb
VITAMIN E	6.81	IU/lb

TFP NUTRITION
NACOGDOCHES, TX

SWEET SHOW LAMB "16"


A medicated corn, oat, and cottonseed hull ration that is specially formulated to promote accelerated growth and weight gain plus optimum finish for show lambs. Also aids in the prevention of urinary calculi and coccidiosis.

FEATURES

- * 1351 Kcal/lb
- * 16% Protein
- * Medicated with Bovatec®
- * Contains cottonseed hulls as a source of fiber
- * Fortified with vitamins and minerals
- * Contains ammonium chloride
- * Fortified with a probiotic that contains:
 1. Encapsulated microbes - Enterococcus faecium & Lactobacillus acidophilus
 2. Yeast culture
 3. Enzymes - Protease, Amylase, and Cellulase

BENEFITS

- * Promotes maximum growth and weight gain.
- * Corn and pearled barley provide optimum finish.
- * Increased average daily gain.
- * Decreased cost/lb of gain.
- * Vitamins and minerals for balanced nutrition.
- * Aids in the prevention of urinary calculi and coccidiosis.
- * Added microbes colonize intestine with protective microflora resulting in better health.
- * Yeast culture improves palatability, digestion, and nutrient utilization.
- * Added enzymes improve digestion of protein, starches, and cellulose resulting in improved feed conversion.
- * Roasted soybeans provide fat and rumen by-pass protein.



For the prevention of coccidiosis caused by *Eimeria ovina*, *Eimeria crandallii*, *Eimeria ovinoidalis* (*Eimeria ninakohlyakimovae*), *Eimeria parva* and *Eimeria intricata* in sheep maintained in confinement.

ACTIVE DRUG INGREDIENT: Lasalocid 20 gram/ton

GUARANTEED ANALYSIS

Crude Protein, minimum	16.00%
(This includes not more than 1.00% equivalent crude protein from non-protein nitrogen.)	
Crude Fat, minimum	4.50%
Crude Fiber, maximum	9.00%
Calcium (Ca), minimum	0.90%
Calcium (Ca), maximum	1.25%
Phosphorus (P), minimum	0.40%
Salt (NaCl), minimum	0.50%
Salt (NaCl), maximum	1.00%
Selenium (Se), minimum	0.20 ppm
Vitamin A, minimum	5,000 IU/LB

INGREDIENTS

Chopped corn, oats, soybean meal, wheat middlings, cane molasses, roughage products (9%), roasted soybeans, pearled barley, flaked corn, rice bran, vegetable oil, calcium carbonate, dicalcium phosphate, ammonium chloride, salt, yeast culture, dried *Lactobacillus acidophilus* fermentation product, dried *Enterococcus faecium* fermentation product, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, ferrous sulfate, ethylenediamine dihydroiodide, cobalt carbonate, natural and artificial flavors.

FEEDING DIRECTIONS

Depending upon body weight, feed continuously to lambs at the rate of 1.5 lb to 7 lb per head per day. This will provide not less than 15 mg nor more than 70 mg of Lasalocid per head per day. Begin feeding immediately after weaning and continue throughout the show season. Additional roughage may be fed if desired to control weight gains.

IMPORTANT MANAGEMENT PRACTICES

1. Changes in the feeding program should be made gradually. Changes should not exceed 1/2 pound per head per day.
2. Provide plain salt free choice.
3. Feed only the amount of feed lambs will clean up in one day. Do not allow old feed to accumulate in feeders.
4. Lambs should be vaccinated against enterotoxemia.
5. Consult your local veterinarian for a recommended health program.

CAUTION: The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. Use only as directed. Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.