



For Improved Digestion, Health, Appearance, and Overall Performance

SHOW FEEDS

PHODUCI	DAIA	
DIG. ENERGY	1325.7	2 Kcal/lb
ME	1.09	Mcal/lb
NE MAINTENANCE	0.73	Mcal/lb
NE GAIN	0.49	Mcal/lb
TDN	66.22	%
PROTEIN	18.17	%
DIG. PROTEIN	15.35	%
NPN	0.80	%
CELLULOSE	6.38	%
FAT	2.62	%
FIBER	7.88	%
LIGNIN	2.26	%
CALCIUM	1.13	%
CHLORINE	0.53	%
COBALT	0.27	mg/lb
IODINE	0.23	mg/lb
IRON		mg/lb
MAGNESIUM	0.21	%
MANGANESE	38.42	mg/lb
PHOSPHORUS	0.53	%
POTASSIUM	1.04	%
SELENIUM	0.09	mg/lb
SODIUM	0.34	%
SULFUR	0.22	%
ZINC	36.18	mg/lb
CAROTENE	0.33	mg/lb
VITAMIN A	5.06	KIU/lb
VITAMIN D	1.23	KIU/lb
VITAMIN E	7.07	IU/lb

CAUTION: Changes in the feeding rate should be made gradually. Feed should be stored in a wellventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.

TEXAS FARM PRODUCTS COMPANY

NACOGDOCHES, TX

SWEET SHOW LAMB "18" MEDICATED

A medicated corn, oat, and cottonseed hull ration that is specially formulated to promote accelerated growth and weight gain plus optimum finish for show lambs. Also aids in the prevention of urinary calculi and coccidiosis.


FEATURES

- 1326 Kcal/lb
- 18% Protein
- Medicated with BovateC®
- Contains cottonseed hulls as a source of fiber
- Fortified with vitamins and minerals
- Contains ammonium chloride
- Contains NO urea or copper sulfate
- Fortified with a probiotic that contains:
- 1. Encapsulated microbes -Streptococcus faecium & Lactobacillus acidophilus
- 2. Yeast culture
- 3. Enzymes Protease, Amylase, and Cellulase

BENEFITS

- * Promotes maximum growth and weight gain.
- Corn provides optimum finish.
- Increased average daily gain.
- Decreased cost/lb of gain.
- Vitamins and minerals for balanced nutrition.
- * Aids in the prevention of urinary calculi and coccidiosis.
- * Pelleted for easy consumption with little waste.
- * Added microbes colonize intestine with protective microflora resulting in better health.
- * Yeast culture improves palatability, digestion, and nutrient utilization.
- Added enzymes improve digestion of protein, starches, and cellulose resulting in improved feed conversion.

Medicated





For the prevention of coccidiosis caused by INGREDIENTS Eimeria ovina, Eimeria crandallis, Eimeria Grain products, plant protein products, ovinoidalis (Eimeria ninakohlyakimovae), Eimeria parva and Eimeria intricata in sheep maintained in confinement.

ACTIVE DRUG INGREDIENT

Lasalocid	20 gram/ton
GUARANTEED ANALYSIS	
Omisto Doubeto metalismos	40.000/

Crude Protein, minimum 18.00% (This includes not more than 0.85% equivalent crude protein from non-protein nitrogen.)

orado protoni nom mon protoni introgom	,
Crude Fat, minimum	2.50%
Crude Fiber, maximum	9.00%
Calcium (Ca), minimum	0.90%
Calcium (Ca), maximum	1.25%
Phosphorus (P), minimum	0.40%
Salt (NaCl), minimum	0.50%
Salt (NaCl), maximum	1.00%
Selenium (Se), minimum 0.2	20 ppm
Vitamin A, minimum 5,000) IU/LB

processed grain by-products, animal protein products, roughage products, cane molasses, dicalcium phosphate, calcium carbonate, salt, ammonium chloride, Streptococcus faecium fermentation product, dried Lactobacillus acidophilus fermentation product, dried Aspergillus oryzae fermentation extract, dried Aspergillus niger fermentation extract, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, ferrous sulfate, ethylenediamine dihydroiodide, cobalt carbonate.

FEEDING DIRECTIONS

Depending upon body weight, feed continuously to lambs at the rate of 1.5 lb to 7 lb per head per day. This will provide not less than 15 mg nor more than 70 mg of Lasalocid per head per day. Begin feeding immediately after weaning and continue throughout the show season. Additional roughage may be fed if desired to control weight gains.

IMPORTANT MANAGEMENT PRACTICES

- 1. Changes in the feeding program should be made gradually. Changes should not exceed 1/2 pound per head per day.
- 2. Provide plain salt free choice.
- 3. Feed only the amount of feed lambs will clean up in one day. Do not allow old feed to accumulate
- 4. Lambs should be vaccinated against enterotoxemia.
- 5. Consult your local veterinarian for a recommended health program.

CAUTION: The safety of Lasalocid in unapproved species has not been established. Use only as directed.