



**Medicated**

**GOAT FEEDS**

**1676**

**PRODUCT DATA**

DIG. ENERGY .....	1125.33	Kcal/lb
ME .....	0.92	Mcal/lb
NE MAINTENANCE ..	0.59	Mcal/lb
NE GAIN .....	0.39	Mcal/lb
TDN .....	55.94	%
PROTEIN .....	15.12	%
DIG. PROTEIN .....	11.62	%
NPN .....	1.20	%
CELLULOSE .....	10.24	%
FAT .....	2.79	%
FIBER .....	15.92	%
LIGNIN .....	4.90	%
CALCIUM .....	1.25	%
CHLORINE .....	0.38	%
COBALT .....	0.25	mg/lb
COPPER .....	5.25	mg/lb
IODINE .....	0.17	mg/lb
IRON .....	55.74	mg/lb
MAGNESIUM .....	0.45	%
MANGANESE .....	71.40	mg/lb
PHOSPHORUS .....	0.54	%
POTASSIUM .....	0.98	%
SELENIUM .....	0.14	mg/lb
SODIUM .....	0.35	%
SULFUR .....	0.18	%
ZINC .....	42.38	mg/lb
CAROTENE .....	4.54	mg/lb
VITAMIN A .....	6.69	KIU/lb
VITAMIN D .....	1.23	KIU/lb
VITAMIN E .....	11.71	IU/lb

TEXAS FARM PRODUCTS COMPANY  
NACOGDOCHES, TX

**COMMERCIAL GOAT PELLETS**


An economical, medicated ration formulated for meat producing goats.

**FEATURES**

- \* 1125 Kcal/lb
- \* 15% Protein
- \* Medicated with Decoquinatate
- \* Ammonium Chloride
- \* Fortified with Vitamins and Minerals

**BENEFITS**

- \* Promotes growth and weight gains.
- \* Medication aids in prevention of coccidiosis in young goats.
- \* Ammonium chloride helps to prevent urinary calculi.
- \* Easy to feed 1/8" diameter pellets ensure uniform consumption.



**ACTIVE DRUG INGREDIENT**

Decoquinatate ..... 27.2 grams/ton

**GUARANTEED ANALYSIS**

Crude Protein, minimum .....	15.00%
(This includes not more than 1.2% equivalent crude protein from non-protein nitrogen.)	
Crude Fat, minimum .....	2.50%
Crude Fiber, maximum .....	16.00%
Calcium (Ca), minimum .....	0.95%
Calcium (Ca), maximum .....	1.25%
Phosphorus (P), minimum .....	0.40%
Salt (NaCl), minimum .....	0.25%
Salt (NaCl), maximum .....	0.75%
Copper (Cu), minimum .....	10 ppm
Copper (Cu), maximum .....	20 ppm
Selenium (Se), minimum .....	0.1 ppm
Vitamin A, minimum .....	5,000 IU/LB

**INGREDIENTS**

Processed grain by-products, roughage products (20%), grain products, plant protein products, forage products, cane molasses, calcium carbonate, diatomaceous earth, ammonium chloride, salt, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, ethylenediamine dihydroiodide, cobalt carbonate.

**FEEDING DIRECTIONS**

Feed according to the chart below to provide 22.7 mg of decoquinatate per 100 pounds of body weight per day.

<b>Weight of Goat (Pounds)</b>	<b>Pounds of Commercial Goat Pellets/Day</b>
20	0.33
40	0.66
60	1.00
80	1.33
100	1.67

Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that coccidiosis is likely to be a hazard. Provide plenty of good quality hay or grazing free choice along with a supply of fresh, clean water.

**WARNING:** Do not feed to goats producing milk for food.

**CAUTION:** Changes in the feeding rate should be made gradually. Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.