



For Improved Digestion, Health, Appearance, and Overall Performance

Medicated

Flavor

Enhanced

# 1701

#### PRODUCT DATA

| PRODUCT          | DATA      |         |
|------------------|-----------|---------|
| DIG. ENERGY      | . 1203.11 | Kcal/lb |
| ME               | 0.98      | Mcal/lb |
| NE MAINTENANCE . | 0.62      | Mcal/lb |
| NE GAIN          | 0.39      | Mcal/lb |
| TDN              | . 59.97   | %       |
| PROTEIN          |           | %       |
| DIG. PROTEIN     | . 12.27   | %       |
| NPN              | 1.20      | %       |
| CELLULOSE        | . 17.84   | %       |
| FAT              | 2.21      | %       |
| FIBER            | . 18.69   | %       |
| LIGNIN           | 5.44      | %       |
| CALCIUM          | 1.00      | %       |
| CHLORINE         | 0.71      | %       |
| COBALT           | 0.40      | mg/lb   |
| COPPER           | 5.33      | mg/lb   |
| IODINE           | 0.30      | mg/lb   |
| IRON             | . 118.17  | mg/lb   |
| MAGNESIUM        | 0.23      | %       |
| MANGANESE        | 45.35     | mg/lb   |
| PHOSPHORUS       | 0.43      | %       |
| POTASSIUM        | 1.32      | %       |
| SELENIUM         | 0.12      | mg/lb   |
| SODIUM           | 0.57      | %       |
| SULFUR           | 0.18      | %       |
| ZINC             | . 59.01   | mg/lb   |
| CAROTENE         | . 15.39   | mg/lb   |
| VITAMIN A        | . 16.15   | KIU/lb  |
| VITAMIN D        | 2.46      | KIU/lb  |
| VITAMIN E        | 21.60     | IU/lb   |
|                  |           |         |

TEXAS FARM PRODUCTS COMPANY NACOGDOCHES. TX

# SHOW FEEDS

## **GOAT GROW & SHOW MEDICATED**

A medicated, alfalfa, corn, and soybean hull based ration formulated to provide the extra nutrition needed to produce champion show goats.

#### **FEATURES** 1203 Kcal/lb \* 17% Protein \* Alfalfa Soybean Hulls & Cottonseed Hulls Fish Meal Ammonium Chloride

- \*Fortified with Probiotics
- Medicated with
- Decoquinate

#### **BENEFITS**

- \* Promotes maximum growth and weight gains
- Energy/Fiber ration allow flexibility in feeding rates
- \* Ammonium chloride helps prevent formation of urinary calculi
- \* Fish meal provides a source of bypass protein
- \* Probiotics improve digestion, health, and overall performance
- \* Medication aids in prevention of coccidiosis in young goats
- Easy to feed 1/8" diameter pellets ensure uniform consumption

For the prevention of coccidiosis in young goats INGREDIENTS caused by Eimeria christenseni and Eimeria Dehydrated alfalfa meal, ground corn, ninakahlvakima

soybean hulls (15%), cottonseed meal, wheat middlings, cottonseed hulls (7.5%), rice bran, fish meal, cane molasses, calcium carbonate, salt, ammonium chloride, dried Streptococcus faecium fermentation product, dried Lactobacillus acidophilus fermentation product, yeast culture, dried Aspergillus oryzae fermentation extract, dried Aspergillus niger fermentation extract, vitamin A acetate, vitamin D3 supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, ethylenediamine dihydroiodide, cobalt carbonate.

### FEEDING DIRECTIONS

Feed according to the chart below to provide 22.7 mg of decoquinate per 100 pounds of body weight per day.

| Weight of Goat<br>(Pounds)            | Pounds of Goat<br>Grow & Show/Day |  |
|---------------------------------------|-----------------------------------|--|
| , , , , , , , , , , , , , , , , , , , |                                   |  |
| 20                                    | 0.33                              |  |
| 40                                    | 0.66                              |  |
| 60                                    | 1.00                              |  |
| 80                                    | 1.33                              |  |
| 100                                   | 1.67                              |  |

Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that coccidiosis is likely to be a hazard. Provide plenty of good quality hay or grazing free choice along with a supply of fresh, clean water.

WARNING: Do not feed to goats producing milk for food.

CAUTION: For Ruminants Only. Do not feed to horses, mules, colts, ponies, or hogs.

Changes in the feeding rate should be made gradually. Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insectinfested feed to animals as it may cause illness or death.