

1/8" PELLETS

FOR RUMINANTS ONLY

CATTLE FEEDS

0777

PRODUCT DATA

PRODUCT DATA			
DIG. ENERGY 1465.48Kcal/lb			
ME	1.38	Mcal/lb	
NE MAINTENANCE	0.79	Mcal/lb	
NE GAIN	0.50	Mcal/lb	
TDN	73.16	%	
PROTEIN	12.05	%	
DIG. PROTEIN	10.39	%	
NPN	1.72	%	
LYSINE	0.44	%	
FAT	2.12	%	
FIBER	13.76	%	
ASH	3.85	%	
CELL WALLS	30.17	%	
ADF	20.66	%	
CELLULOSE	12.67	%	
LIGNIN	5.74	%	
CALCIUM	0.61	%	
CHLORINE	1.04	%	
COBALT	0.32	mg/lb	
COPPER	8.72	mg/lb	
IRON	61.64	mg/lb	
IODINE	0.27	mg/lb	
MAGNESIUM	0.23	%	
MANGANESE	32.28	mg/lb	
PHOSPHORUS	0.31	%	
POTASSIUM	1.14	%	
SODIUM	0.44	%	
SELENIUM	0.04	mg/lb	
SULFUR	0.22	%	
ZINC		mg/lb	
RIBOFLAVIN	0.76	mg/lb	
THIAMINE	0.92	mg/lb	
NIACIN		mg/lb	
PAN ACID		mg/lb	
CHOLINE		mg/lb	
PYRIDOXINE	3.53	mg/lb	
FOLIC ACID	0.16	mg/lb	
VITAMIN B12	0.00	mcg/lb	
VITAMIN E	6.97	IU/lb	
VITAMIN A		KIU/lb	
VITAMIN D	1.23	KIU/lb	

TFP NUTRITION NACOGDOCHES, TEXAS

BULL & HEIFER DEVELOPER

A cottonseed hull and corn based ration used by many cattlemen raising purebred bulls and heifers for replacement or sale.

FEATURES

- * 1465 kcal/lb
- * 12% protein
- * 21% cottonseed hulls
- * 45% corn
- * Contains soybean hulls

BENEFITS

- *Corn provides optimum finish on animals.
- * Medication improves rate of weight gain.
- * Excellent for fitting cattle for show.
- * Cottonseed hulls and soybean hulls provide high quality roughage.



GUARANTEED ANALYSIS

GUARAN I EED ANALTSIS		
Crude Protein, minimum	12.00%	
(This includes not more than 1.72% eq	uivalent	
crude protein from non-protein nitrogen.)		
Crude Fat, minimum	2.00%	
Crude Fiber, maximum	18.00%	
Calcium (Ca), minimum	0.45%	
Calcium (Ca), maximum	0.75%	
Phosphorus (P), minimum	0.30%	
Salt (NaCl), minimum	0.75%	
Salt (NaCl), maximum	1.25%	

 Potassium (K), minimum
 0.50%

 Vitamin A, minimum
 5,000 IU/Lb

 Vitamin D3, minimum
 1,230 IU/Lb

INGREDIENTS

Grain products, plant protein products, roughage products (28%), cane molasses, calcium carbonate, salt, urea, dicalcium phosphate, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, ethylenediamine dihydroiodide, cobalt carbonate.

FEEDING DIRECTIONS

Feed continuously to beef cattle on pasture at the rate of 4.5 to 15 pounds per head per day. This will provide a minimum of 60 mg and a maximum of 200 mg of lasalocid per head per day. Provide a source of fresh, clean water at all times.

CAUTION: Changes in feed should be made gradually. Feed should be stored in a well-ventilated, dry area that is protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed as it may cause illness or death.