



MINERALS



CATTLE FEEDS

1064

PRODUCT DATA

DIG.ENERGY	173.14	Kcal/lb
ME	0.14	Mcal/lb
NE MAINTENANCE ..	0.10	Mcal/lb
NE GAIN	0.06	Mcal/lb
TDN	8.64	%
PROTEIN	3.76	%
DIG. PROTEIN	2.98	%
NPN	0.00	%
LYSINE	0.15	%
FAT	0.20	%
FIBER	1.18	%
ASH	0.84	%
CELL WALLS	2.52	%
ADF	1.53	%
CELLULOSE	1.08	%
LIGNIN	0.54	%
CALCIUM	13.82	%
CHLORINE	12.93	%
COBALT	14.05	mg/lb
COPPER	909.90	mg/lb
IRON	1250.85	mg/lb
IODINE	36.33	mg/lb
MAGNESIUM	1.57	%
MANGANESE	1881.08	mg/lb
PHOSPHORUS	6.52	%
POTASSIUM	2.32	%
SODIUM	7.99	%
SELENIUM	11.35	mg/lb
SULFUR	0.81	%
ZINC	3647.50	mg/lb
RIBOFLAVIN	0.23	mg/lb
THIAMINE	0.28	mg/lb
NIACIN	2.01	mg/lb
PAN ACID	1.10	mg/lb
CHOLINE	127.00	mg/lb
PYRIDOXINE	0.26	mg/lb
FOLIC ACID	0.09	mg/lb
VITAMIN B12	0.00	mcg/lb
VITAMIN E	80.64	IU/lb
VITAMIN A	95.04	KIU/lb
VITAMIN D	30.01	KIU/lb

TEXAS FARM PRODUCTS COMPANY
NACOGDOCHES, TEXAS

KAL-PHOS 14-6.5 MINERAL SUPPLEMENT WITH BOVATEC®

A self-limiting pasture supplement designed to supply the nutrients most often deficient in forages.

FEATURES

- * High levels of calcium and phosphorus in a 2:1 ratio
- * 20% salt
- * Fortified with vitamins
- * Medicated with Bovatec®

BENEFITS

- * Fewer breeding problems.
- * Healthier calves at birth.
- * Fewer calving problems.
- * Better utilization of poor quality forages.

For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Lasalocid 1440 grams/ton

GUARANTEED ANALYSIS

Calcium (Ca), minimum 13.00%
 Calcium (Ca), maximum 15.00%
 Phosphorus (P), minimum 6.50%
 Salt (NaCl), minimum 19.00%
 Salt (NaCl), maximum 22.00%

INGREDIENTS

Deflourinated phosphate, salt, calcium carbonate, cottonseed meal, potassium chloride, cane molasses, magnesium sulfate, magnesium oxide, potassium sulfate, iron oxide, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, copper sulfate, ethylenediamine dihydroiodide, cobalt sulfate.

FEEDING DIRECTIONS

Feed continuously on a free choice basis to pasture cattle to provide 60-200 mg of lasalocid per head daily. Pasture and roughage should be adequate to assure a minimum consumption of 1.34 oz/head/day (which provides 60 mg of lasalocid) and a maximum consumption of 4.44 oz/head/day (which provides 200 mg of lasalocid). If cattle consume more or less than these levels, the feeders should be moved further or closer to the general resting or watering area. If cattle still fail to consume the correct amount, discontinue feeding this product.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

CAUTION: Feed should be stored in a well-ventilated, dry area that is protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.

