



37% RANGE BLOCK

For Mature Beef Cattle on Pasture

GUARANTEED ANALYSIS:

Crude Protein (Min)	37.00%
(This includes not more than 23.00% equivalent protein from non-protein nitrogen).	
Crude Fat (Min)	0.50%
Crude Fiber (Max)	7.00%
Calcium (Min)	6.00%
Calcium (Max).....	7.00%
Phosphorus (Min).....	0.70%
Salt (Min).....	21.00%
Salt (Max).....	25.00%
Magnesium (Min)	1.00%
Potassium (Min)	1.00%
Copper (Min)	130 ppm
Iodine (Min).....	10 ppm
Manganese (Min)	300 ppm
Selenium (Min)	6 ppm
Zinc (Min).....	500 ppm
Vitamin A (Min)	30,000 IU/lb
Vitamin D-3 (Min).....	7,500 IU/lb
Vitamin E (Min)	7.5 IU/lb

INGREDIENT STATEMENT:

Salt, Plant Protein Products, Calcium Carbonate, Molasses Products, Urea, Magnesium Limestone, Calcium Hydroxide, Dicalcium Phosphate, Monocalcium Phosphate, Potassium Sulfate, Magnesium Sulfate, Manganese Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Chloride, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement and Vitamin E Supplement.

NET WEIGHT: 33.3 LB (15.12 KG)



INTRODUCTORY PERIOD:

Following a 7 to 10 day acclimation period, average consumption should be 1 to 2 pounds per head per day for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

FEEDING RECOMMENDATIONS:

1. Provide 1 block per 5 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding – 10 feet minimum.
3. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two-thirds consumed, provide a new block near it. Place very small pieces in a feed trough to be cleaned up.
4. Cattle should consume 1 to 2 pounds per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate free choice mineral supplement in addition to this supplement.

CAUTION: USE AS DIRECTED. Do not feed to horses. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Batch #

Manufactured For:
TFP NUTRITION

Nacogdoches, Texas 75963-0009

11174