## 24/16 Beef Tub

# Molasses Block SIAR Supplement for Beef Cattle on Pasture

### **GUARANTEED ANALYSIS**

| Crude Protein, minimum                                       | 24.00%    |
|--|-----------|
| (This includes not more than 16.00% equivalent crude protein |           |
| from non-protein nitrogen.)                                  |           |
| Crude Fat, minimum   | 4.00%     |
| Crude Fiber, maximum   | 5.00%     |
| Calcium(Ca), minimum   | 3.00%     |
| Calcium (Ca), maximum  | 4.00%     |
| Phosphorus (P), minimum                                      | 0.50%     |
| Salt (NaCI), minimum   | 5.50%     |
| Salt (NaCl), maximum   | 6.50%     |
| Magnesium (Mg), minimum                                      | 2.00%     |
| Potassium (K), minimum                                       | 0.50%     |
| Copper (Cu), minimum   | 30 ppm    |
|  | 1.0 ppm   |
| Zinc, (Zn), minimum  | 75 ppm    |
| Manganese (Mn), minimum                                      | 55 ppm    |
| lodine (I), minimum  | 2.1 ppm   |
| Cobalt (Co), minimum   | 0.2 ppm   |
| Vitamin A, minimum   | 000 IU/lb |
|  | 000 IU/lb |
| Vitamin E, minimum   | 4 IU/lb   |

#### INGREDIENTS

Molasses products, plant protein products, processed grain byproducts, hydrolyzed vegetable oil, condensed fermented corn extractives,urea, magnesium oxide, salt, calcium oxide, ground limestone, phosphoric acid, sulfuric acid, lignin sulfate, ammonium sulfate, vitamin A supplement, vitamin D<sub>3</sub> supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, copper sulfate, ethylenediamine dihydroiodide, and cobalt sulfate.

### **FEEDING DIRECTIONS**

Feed free choice as a supplement to mature beef cattle grazing adequate forage. Provide a source of fresh, clean water at all times along with Lone Star Kal-Phos 12-12 Mineral Supplement. DO NOT feed this block in conjunction with other feeds containing non-protein nitrogen. Consumption of this block should be approximately 1 to 2 pounds per head per day. The following guidelines should be followed to ensure adequate consumption:

- Provide one 200 pound block for every 10 to 13 head of cattle.
- Place blocks near water source or loafing areas. Make blocks available in more than one location in large pastures. If consumption is too high, move the blocks farther away from the water source and loafing areas, reduce the number of blocks if necessary to control consumption.
- Maintain a consistent supply of blocks. Do not feed to deprived animals as overconsumption may result.

**CAUTION:** USE AS DIRECTED. For ruminants only. Do not feed to horses, mules, colts, ponies, or hogs. Store in a dry area.

Manufactured for: TFP Nutrition Nacogdoches, TX 75963-0009 **2187**