



Medicated

GOAT FEEDS

1676

COMMERCIAL GOAT PELLETS

An economical, medicated ration formulated for meat producing goats.

PRODUCT DATA

DIG. ENERGY	1125.33	Kcal/lb
ME	0.92	Mcal/lb
NE MAINTENANCE ..	0.59	Mcal/lb
NE GAIN	0.39	Mcal/lb
TDN	55.94	%
PROTEIN	15.12	%
DIG. PROTEIN	11.62	%
NPN	1.20	%
CELLULOSE	10.24	%
FAT	2.79	%
FIBER	15.92	%
LIGNIN	4.90	%
CALCIUM	1.25	%
CHLORINE	0.38	%
COBALT	0.25	mg/lb
COPPER	5.25	mg/lb
IODINE	0.17	mg/lb
IRON	55.74	mg/lb
MAGNESIUM	0.45	%
MANGANESE	71.40	mg/lb
PHOSPHORUS	0.54	%
POTASSIUM	0.98	%
SELENIUM	0.14	mg/lb
SODIUM	0.35	%
SULFUR	0.18	%
ZINC	42.38	mg/lb
CAROTENE	4.54	mg/lb
VITAMIN A	6.69	KIU/lb
VITAMIN D	1.23	KIU/lb
VITAMIN E	11.71	IU/lb

TFP NUTRITION
NACOGDOCHES, TX

FEATURES

- * 1125 Kcal/lb
- * 15% Protein
- * Medicated with Decoquinatate
- * Ammonium Chloride
- * Fortified with Vitamins and Minerals

BENEFITS

- * Promotes growth and weight gains.
- * Medication aids in prevention of coccidiosis in young goats.
- * Ammonium chloride helps to prevent urinary calculi.
- * Easy to feed 1/8" diameter pellets ensure uniform consumption.

For the prevention of coccidiosis in young goats caused by *Eimeria christenseni* and *Eimeria ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Decoquinatate 27.2 grams/ton

GUARANTEED ANALYSIS

Crude Protein, minimum	15.00%
(This includes not more than 1.2% equivalent crude protein from non-protein nitrogen.)	
Crude Fat, minimum	2.50%
Crude Fiber, maximum	16.00%
Calcium (Ca), minimum	0.95%
Calcium (Ca), maximum	1.25%
Phosphorus (P), minimum	0.40%
Salt (NaCl), minimum	0.25%
Salt (NaCl), maximum	0.75%
Copper (Cu), minimum	10 ppm
Copper (Cu), maximum	20 ppm
Selenium (Se), minimum	0.1 ppm
Vitamin A, minimum	5,000 IU/LB

FEEDING DIRECTIONS

Feed according to the chart below to provide 22.7 mg of decoquinatate per 100 pounds of body weight per day.

Weight of Goat (Pounds)	Pounds of Commercial Goat Pellets/Day
20	0.33
40	0.66
60	1.00
80	1.33
100	1.67

Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that coccidiosis is likely to be a hazard. Provide plenty of good quality hay or grazing free choice along with a supply of fresh, clean water.

WARNING: Do not feed to goats producing milk for human consumption.

CAUTION: For ruminants only. Do not feed to horse, mules, colts, ponies or hogs. Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed as it may cause illness or death.

