



1/8" PELLETS

FOR RUMINANTS
ONLY

CATTLE FEEDS

0777

PRODUCT DATA

DIG. ENERGY	1477.71	Kcal/lb
ME	1.39	Mcal/lb
NE MAINTENANCE ..	0.79	Mcal/lb
NE GAIN	0.51	Mcal/lb
TDN	73.75	%
PROTEIN	12.06	%
DIG. PROTEIN	10.59	%
NPN	1.72	%
LYSINE	0.43	%
FAT	2.99	%
FIBER	14.47	%
ASH	3.46	%
CELL WALLS	32.34	%
ADF	21.80	%
CELLULOSE	13.21	%
LIGNIN	5.90	%
CALCIUM	0.57	%
CHLORINE	0.77	%
COBALT	0.27	mg/lb
COPPER	6.01	mg/lb
IRON	57.69	mg/lb
IODINE	0.18	mg/lb
MAGNESIUM	0.26	%
MANGANESE	43.94	mg/lb
PHOSPHORUS	0.40	%
POTASSIUM	0.91	%
SODIUM	0.43	%
SELENIUM	0.05	mg/lb
SULFUR	0.26	%
ZINC	33.77	mg/lb
RIBOFLAVIN	0.68	mg/lb
THIAMINE	1.61	mg/lb
NIACIN	17.80	mg/lb
PAN ACID	3.44	mg/lb
CHOLINE	318.68	mg/lb
PYRIDOXINE	3.75	mg/lb
FOLIC ACID	0.15	mg/lb
VITAMIN B12	0.00	mcg/lb
VITAMIN E	9.53	IU/lb
VITAMIN A	5.54	KIU/lb
VITAMIN D	1.23	KIU/lb

TFP NUTRITION
NACOGDOCHES, TEXAS

BULL & HEIFER DEVELOPER

A cottonseed hull and corn based ration used by many cattlemen raising purebred bulls and heifers for replacement or sale.

FEATURES

- * 1477 kcal/lb
- * 12% protein
- * 21% cottonseed hulls
- * 45% corn
- * Contains soybean hulls

BENEFITS

- * Corn provides optimum finish on animals.
- * Excellent rate of weight gain.
- * Excellent for fitting cattle for show.
- * Cottonseed hulls and soybean hulls provide high quality roughage.

GUARANTEED ANALYSIS

Crude Protein, minimum	12.00%
(This includes not more than 1.72% equivalent crude protein from non-protein nitrogen.)	
Crude Fat, minimum	2.00%
Crude Fiber, maximum	18.00%
Calcium (Ca), minimum	0.45%
Calcium (Ca), maximum	0.75%
Phosphorus (P), minimum	0.30%
Salt (NaCl), minimum	0.75%
Salt (NaCl), maximum	1.25%
Potassium (K), minimum	0.50%
Vitamin A, minimum	5,000 IU/Lb
Vitamin D3, minimum	1,230 IU/Lb

INGREDIENTS

Grain products, plant protein products, roughage products (28%), cane molasses, calcium carbonate, salt, urea, dicalcium phosphate, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, ethylenediamine dihydroiodide, cobalt carbonate.

FEEDING DIRECTIONS

Feed to beef cattle on pasture at the rate of 5 to 15 pounds per head per day. Provide a source of fresh, clean water at all times.

CAUTION: Use only as directed. Do not feed to horses, mules, colts, ponies or hogs. Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed to animals as it may cause illness or death.

